

AURYXIA—The first iron tablet *specifically made with your kidneys in mind*



WHAT IS IRON DEFICIENCY ANEMIA?

- Iron deficiency anemia occurs when the body doesn't have enough iron to make hemoglobin, a protein needed for healthy red blood cells
- Hemoglobin is needed so red blood cells can carry the oxygen your body needs
- Symptoms may include weakness, lack of energy, and shortness of breath



WHAT AURYXIA DOES:

- AURYXIA (ah-RICKS-ee-ah) is a prescription medicine that treats iron deficiency anemia in adults with chronic kidney disease not on dialysis
- AURYXIA can increase hemoglobin and iron levels in the body. The iron in AURYXIA gets digested in the same way as the iron in your food. This allows your body to keep the iron it needs to function properly, and remove what it doesn't need as waste



TAKING AURYXIA:

- AURYXIA does not require needles, infusions centers, or leaving the comfort of your home
- AURYXIA tablets should be swallowed whole, and not chewed or crushed
- The suggested starting dose is 1 tablet, 3 times per day with meals
- You should always take AURYXIA as prescribed by your doctor



YOU MAY NOTICE:

- AURYXIA contains iron and may cause dark stools, which is considered normal when taking oral medications containing iron
- Side effects may include diarrhea, constipation, nausea, abdominal pain, and hyperkalemia (high levels of potassium in the blood)



TAKING AURYXIA WITH OTHER MEDICATIONS?

- **Doxycycline**—Take at least 1 hour before AURYXIA
- **Ciprofloxacin**—Take at least 2 hours before or after AURYXIA
- Be sure to tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements

NOTES

Work with your healthcare provider to reach your hemoglobin goal

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about AURYXIA[®] (ferric citrate)?

AURYXIA contains iron. Keep it away from children to prevent an accidental ingestion of iron and potentially fatal poisoning. Call a poison control center or your healthcare provider if a child swallows AURYXIA.

AURYXIA can increase iron levels in your blood. Iron absorbed from AURYXIA may also increase iron in your body. Your healthcare provider will monitor your iron levels. If you are receiving intravenous (IV) iron, your IV iron dose may be adjusted or discontinued.

Please see full **Important Safety Information** on the following page. The FDA-approved product labeling can be found at [AURYXIA.com](https://www.auryxia.com)



Scan to learn more at AURYXIA.com

IMPORTANT SAFETY INFORMATION

This information does not take the place of talking to your healthcare provider about your medical condition or treatment

What is the most important information I should know about AURYXIA® (ferric citrate)?

AURYXIA contains iron. Keep it away from children to prevent an accidental ingestion of iron and potentially fatal poisoning. Call a poison control center or your healthcare provider if a child swallows AURYXIA

AURYXIA can increase iron levels in your blood. Iron absorbed from AURYXIA may also increase iron in your body. Your healthcare provider will monitor your iron levels. If you are receiving intravenous (IV) iron, your IV iron dose may be adjusted or discontinued

What is AURYXIA?

For adults who have iron deficiency anemia, as well as CKD, and are NOT on dialysis, AURYXIA is a prescription medicine that can increase hemoglobin levels and increase iron levels in the body

Who should not take AURYXIA?

Do not take AURYXIA if you have been diagnosed with an iron overload syndrome, such as hemochromatosis

AURYXIA may not be right for you. Before starting AURYXIA, tell your healthcare provider if you:

- have any other medical conditions
- are pregnant, plan to become pregnant, are breastfeeding or plan to breastfeed

Tell your healthcare provider about all of the medicines you take, including:

- the antibiotics doxycycline or ciprofloxacin
- prescription and over-the-counter medicines, vitamins, and herbal supplements

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine

How should I take AURYXIA?

- Take AURYXIA exactly as prescribed by your healthcare provider
- Take AURYXIA with meals and adhere to any diet prescribed by your healthcare provider
- Your healthcare provider will tell you how much AURYXIA to take and may change your dose if necessary
- Swallow AURYXIA whole. Do not chew or crush.
- If you are taking the antibiotics doxycycline or ciprofloxacin, you will need to take it separately from AURYXIA. Follow your healthcare provider's instructions on when to take doxycycline or ciprofloxacin while you are also taking AURYXIA

What are the most common side effects of AURYXIA?

The most common side effects of AURYXIA for use in the treatment of iron deficiency anemia in adult patients with CKD not on dialysis include:

- diarrhea
- constipation
- nausea
- abdominal pain
- high levels of potassium in the blood

AURYXIA contains iron and may cause dark stools, which are considered normal with oral medications containing iron

These are not all the side effects of AURYXIA. For more information ask your healthcare provider or pharmacist

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Call your healthcare provider for medical advice about side effects. You may report suspected side effects to Akebia Therapeutics, Inc. at 1-844-445-3799 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

How should I store AURYXIA?

- Store AURYXIA between 68 to 77°F (20 to 25°C)
- Keep AURYXIA tablets dry

The FDA-approved product labeling can be found at AURYXIA.com